

Phytochemical Profiling And Pharmacognostical Evaluation Of Ornamental Plant *Rhaphidophora Aurea* (Money Plant)

Mr. Rajat Yadav^{1*}, Dr. Kratika Daniel²

^{1*}Research scholar, Oriental University, Indore M.P. Phone no: 7906575824, Email: yadavm.pharm@gmail.com

²Professor, Oriental University, Indore M.P.

KEYWORDS

Rhaphidophora aurea,
Phytochemical
Screening,
Pharmacognostical
Evaluation,
Standardization
Parameters, Herbal Drug
Quality

ABSTRACT

The present study aims to investigate the phytochemical and pharmacognostical attributes of *Rhaphidophora aurea* (Money Plant), an ornamental plant with potential medicinal properties. A detailed pharmacognostic evaluation was conducted, including macroscopic and microscopic examination, powder analysis, and physicochemical characterization, to establish a comprehensive reference for academic and commercial applications. The phytochemical screening involved methanol (80%), ether, and chloroform extracts, which were analyzed for the presence of secondary metabolites such as alkaloids, glycosides, tannins, proteins, amino acids, and terpenoids. The determination of ash values played a crucial role in standardizing the crude drug and assessing the presence of extraneous matter. Furthermore, qualitative screening for saponins, tannins, alkaloids, terpenoids, anthraquinones, flavonoids, and glycosides was performed to authenticate the medicinal potential of the plant. The study provides critical physicochemical parameters necessary for ensuring the quality and purity of the plant material, thereby contributing to its standardization. These findings serve as a scientific basis for future research and potential applications in herbal medicine, supporting the establishment of pharmacopoeial standards for *Rhaphidophora aurea*.

1. Introduction:

Medicinal plants have been extensively studied for their bioactive compounds, contributing significantly to traditional and modern medicine. Among them, *Rhaphidophora aurea* (commonly known as the Money Plant) is an ornamental plant widely cultivated for its aesthetic appeal and air-purifying properties. However, its phytochemical composition and potential pharmacognostical significance remain underexplored. The phytochemical constituents of medicinal plants are responsible for their therapeutic effects, making phytochemical screening essential for identifying bioactive compounds that may have pharmacognostical relevance (Gurib-Fakim, 2014).

The genus *Rhaphidophora*, belonging to the family Araceae, is known for its diverse phytoconstituents, including flavonoids, alkaloids, tannins, and saponins, which have been reported in related species (Kumar et al., 2020). These secondary metabolites play a crucial role in plant defense and may offer beneficial properties such as antioxidant and antimicrobial activities (Singh et al., 2019). While *Rhaphidophora aurea* is primarily recognized for its ornamental value, its potential medicinal attributes remain largely unexplored. Hence, comprehensive phytochemical profiling and pharmacognostical evaluation are imperative to establish its therapeutic significance.

Pharmacognostical evaluation involves a detailed investigation of a plant's macroscopic, microscopic, physicochemical, and phytochemical characteristics. Standardization parameters such as moisture content, extractive values, ash values, and fluorescence analysis provide essential quality control data, ensuring consistency in herbal formulations (Kokate et al., 2019). Additionally, microscopic analysis of leaf structures, stomatal arrangement, and trichome morphology aids in plant identification and authentication, preventing adulteration and substitution in herbal medicine (Trease & Evans, 2016).

Given the increasing global interest in natural products and herbal therapeutics, this study aims to perform a detailed phytochemical screening and pharmacognostical evaluation of *Rhaphidophora aurea*. This research will contribute to the standardization of this species and provide scientific validation for its potential inclusion in herbal medicine formulations.

2. Material and Methodology:

2.1. Collection and Plant Authentication

The leaf of the climber was possessed from the herbal plant garden of Shri Ram Murti Smarak College of Engineering and Technology Pharmacy, Bareilly, Uttar Pradesh (India). The plant authentication was done by Dr. Alok Kumar Shrivastav, Department of Botany, M.J.P. Rohilkhand University,

Bareilly Uttar Pradesh, India and was identified as *Rhaphidophora aurea* (Money Plant) holding reference letter no. RU/PS/2023/07 dated April 5th 2023.

2.2. Macroscopic and Microscopic Evaluation:

2.2.1 Macroscopic Analysis:

The macroscopic characteristics of the leaves, such as shape, size, color, texture, venation pattern, and surface features, were recorded following standard protocols (Kokate et al., 2019).

2.2.2. Microscopic Analysis:

Transverse sections of the fresh leaves were prepared and stained with safranin and iodine-potassium iodide (I2KI) for anatomical studies. The microscopic features, including epidermal cells, stomatal type, trichomes, and vascular bundles, were observed under a compound microscope (Trease & Evans, 2016). Powder microscopy of dried leaf powder was performed using chloral hydrate and phloroglucinol-HCl staining to identify lignified tissues and starch grains.

2.3. SEM Image of leaf surface of the money plant dry powder (MLDP):

The prepared sample was placed in the SEM chamber, and imaging was conducted under high vacuum conditions using an accelerating voltage of 5-20 kV, following standard SEM protocols for plant samples (Bozzola & Russell, 2020). Magnifications ranging from 500x to 5000x were used to capture the detailed microstructure of the leaf powder, focusing on epidermal features, trichomes, and stomatal patterns. Previous studies on SEM imaging of plant powders have demonstrated that high-magnification analysis allows for distinguishing between species based on surface morphology and microstructural characteristics (Medeiros et al., 2021).

2.3. Powder microscopy:

The fresh leaves of *Rhaphidophora aurea* were collected, washed with distilled water, shade-dried at room temperature for 10-12 days, and finely powdered using a mechanical grinder. The powder was then sieved through a 60-mesh sieve to obtain a uniform consistency and stored in an airtight container for analysis. A small amount of powdered leaf sample was mounted on a clean glass slide, and the following staining procedures were performed to detect different cellular components:

- **Lignified Tissues:** A drop of 1% phloroglucinol solution was added to the powder, followed by a few drops of concentrated hydrochloric acid. The slide was observed under a microscope for pink to red-stained lignified elements.
- **Starch Grains:** The powder was treated with iodine solution, and the presence of starch was confirmed by the appearance of blue-black coloration.
- **Calcium Oxalate Crystals:** The powder was mixed with a few drops of glycerin and observed for birefringent crystals under polarized light.
- **Oil Globules:** The powder was stained with Sudan III, which imparted a red color to the oil droplets.
- **Cellulose and Parenchyma Cells:** The powder was stained with safranin, and the structural details of parenchymatous cells and fibers were examined.



Figure 1: Powder Of *Rhaphidophora aurea* Leaves

2.3. Physicochemical evaluation:

2.3.1. Determination of Moisture Content:

Loss on drying (LOD) refers to the percentage (w/w) of weight loss due to evaporation of water and other vapour substances under specific conditions. A sample of 1 gram of dried *Rhaphidophora aurea* is placed in a hollow vial and weighed. The specimen is then uniformly spread and dehydrated in hot air oven at 105°C for one hour with the bung left removed. After this period, the bung is sealed, and the bottle is left to be level-headed to surrounding temperature before being weighed again (Indian Pharmacopoeia, 2021).

2.3.2. Determination of Ash Values:

2.3.2.1. Total Ash Value:

The total material left after incineration of the powdered drug at temperature 400°C to remove all carbon atoms is determined. Accurately weigh 2-3 g of finely powdered *Rhaphidophora aurea* leaf sample. Place the sample into a previously ignited, cooled, and tarred crucible to ensure no pre-existing residue affects the weight measurement (Trease & Evans, 2016). Gradually heat the sample on a low flame using a hot plate until all moisture is removed and the material is charred without flaming. This prevents excessive loss of volatile inorganic constituents. Transfer the crucible to a preheated muffle furnace and incinerate the sample at 500-600°C for 4-6 hours or until a constant weight is obtained. Ensure that the ash is white or grayish-white, indicating complete oxidation of organic matter (Kokate et al., 2019). Remove the crucible from the furnace using tongs and place it in a desiccator to cool to room temperature. Weigh the crucible accurately and record the weight of the ash.

The total ash content is expressed as a percentage using the formula:

$$\text{Total Ash (\%)} = \left(\frac{\text{Weight of Ash Residue}}{\text{Weight of Sample Taken}} \right) \times 100$$



Figure 2: Total Ash

2.3.2.2. Water-soluble ash:

Accurately weigh 2-3 g of finely powdered *Rhaphidophora aurea* leaf sample and transfer it into a previously ignited and tarred crucible. Incinerate the sample in a muffle furnace at 500-600°C until a constant weight is obtained, ensuring complete combustion of organic matter. Cool the crucible in a desiccator and weigh the total ash obtained. Add 25 mL of distilled water to the total ash residue in a 100 mL glass beaker. Stir well and allow the mixture to stand for 10-15 minutes to ensure the complete dissolution of water-soluble inorganic salts (Kokate et al., 2019). Filter the solution through an ash-free filter paper into a pre-weighed sintered crucible or porcelain dish. Rinse the filter paper with hot distilled water to ensure maximum recovery of soluble constituents. The filtrate is evaporated to dryness on a hot plate and further dried at 105°C in an oven until a constant weight is achieved. The dried residue represents the water-soluble ash fraction. Cool the crucible in a desiccator and weigh the final water-insoluble residue.

The percentage of water-soluble ash is calculated using the formula:

$$\text{Water-Soluble Ash (\%)} = \left[\frac{\text{Total Ash Weight} - \text{Water-Insoluble Ash Weight}}{\text{Weight of Sample Taken}} \right] \times 100$$

2.3.2.3. Acid Insoluble Ash:

Acid-insoluble ash represents the portion of total ash that remains undissolved when treated with dilute hydrochloric acid. This fraction primarily includes silica, sand, and other earthy materials that may have been

incorporated into the plant sample during cultivation, handling, or processing (Trease & Evans, 2016). Accurately weigh 2-3 g of finely powdered *Rhaphidophora aurea* leaves and transfer it to a previously ignited and tared crucible. Incinerate the sample in a muffle furnace at 500-600°C until free from carbon (i.e., a white or grayish-white ash is obtained). Cool the crucible in a desiccator and weigh the ash. This represents the total ash value. Moisten the total ash residue with a few drops of distilled water. Add 25 mL of dilute hydrochloric acid (6N) to the crucible and cover it with a watch glass to prevent loss by splashing. Heat the mixture on a hot plate for 5 minutes, stirring occasionally to ensure thorough digestion of soluble ash components. Filter the solution through an ash-free filter paper into a pre-weighed sintered crucible or porcelain dish. The residue collected on the filter paper is washed thoroughly with hot distilled water to remove any soluble impurities. The washed residue is then transferred back to the crucible and ignited in a muffle furnace at 500-600°C for about 1 hour until a constant weight is obtained. The crucible is cooled in a desiccator and weighed. The acid-insoluble ash percentage is determined using the formula:

$$\text{Acid-Insoluble Ash (\%)} = \left[\frac{\text{Weight of acid-Insoluble residue}}{\text{Weight of sample taken}} \right] \times 100$$



Figure 3: Acid-insoluble Ash

2.3.3. Extractive Values:

The extractive value of the herbal drug ascertains the quantity of functioning content extirpated with solvent from a fixed quantity of medicinally potent herb, providing acumen into the present chemical ingredients present.

2.3.3.1. Alcohol-Soluble Extractive Value:

Weigh accurately 5 g of the finely powdered plant material and transfer it into a 250 mL conical flask. Add 100 mL of ethanol to the flask and shake the mixture well. Allow the sample to macerate for 24 hours at room temperature, shaking occasionally to ensure uniform extraction. After 24 hours, filter the extract through Whatman No. 1 filter paper into a pre-weighed evaporating dish to remove undissolved residues. The filtrate is evaporated to dryness using a water bath at 60°C until a constant weight is obtained. The dried extract is cooled in a desiccator and weighed.

The alcohol-soluble extractive value (%) is calculated using the formula:

$$\text{Alcohol-Soluble Extractive (\%)} = \left(\frac{\text{Weight of dried extract}}{\text{Weight of sample taken}} \right) \times 100$$

2.3.3.2. Water-Soluble extractive value:

Weigh accurately 5 g of the finely powdered plant material and transfer it into a 250 mL conical flask. Add 100 mL of distilled water to the flask and shake the mixture well. Allow the sample to macerate for 24 hours at room temperature with occasional shaking. After 24 hours, filter the solution through a Whatman No. 1 filter paper into a pre-weighed evaporating dish to remove insoluble residues. The filtrate is evaporated to dryness in a water bath at 60°C until a constant weight is obtained. The dried extract is cooled in a desiccator and weighed.

The water-soluble extractive value (%) is calculated using the formula:

$$\text{Water-Soluble Extractive (\%)} = \left(\frac{\text{Weight of dried extract}}{\text{Weight of sample taken}} \right) \times 100$$

2.4. Extract Preparation:

The Leaves of the plant were segregated, washed thoroughly with common water to make them free from dust particles and other unwanted materials accumulated on leaves from environment, leaves of the plant were shade

dried for 10-12 days (23–27°C) and homogenized to the plant material was ground into a coarse powder using a grinder. After grinding, the powdered materials were stored in a plastic bag for extraction. The coarse powder was then filtered through an 80-mesh sieve. A total of 250 g of powdered *Rhabdophora aurea* plant material was extracted by using petroleum ether, chloroform, and 80% methanol with the help of Soxhlet apparatus. All the obtained extracts were subsequently analyzed for the presence of phytoconstituents to determine their composition.

2.5. Phytochemical Screening:

2.5.1. Alkaloid Detection assay:

Mayer's Test: 2 ml portion of the liquid was mixed with 2N HCl, and the resulting water-fusible layer being decanted. Potassium mercuric Iodide solution was then mixed to this layer. The appearance of a creamy-colored ppt indicated and confirmed the existence of alkaloids.

Dragendroff's Test: Similar being to Mayer's assay, 2 ml portion of liquid herbal decant being mixed with Dragendroff's reagent. The appearance of reddish-brown precipitate with Potassium bismuth iodide suggested alkaloid content.

Wagner's Test: Another method involved mixing a 2 ml portion of the liquid with Iodo Potassium iodide reagent. The appearance of a reddish-brown settled ppt. confirmed alkaloid.

Hager's Test: 2 millilitres portioned liquid was combined with Picric acid reagent. The developed yellow precipitate marked alkaloid content.



Figure 4: Alkaloid Detection

2.5.2. Glycoside Detection Test:

Fehling's Solution Test: A 2 ml portion of the liquid was mixed with Fehling's Solution A and Fehling's Solution B in equal amounts, the mixture was warmed up. The appearance of a brick-red colored precipitate marked existence of glycosides.

Legal's examination: The sample was mixed with a 2 ml solution to which basic sodium nitroprusside and pyridine was added, resulting in a blood red color.

2.5.3. Flavonoid detection test:

Shinoda method: A few pieces of magnesium ribbon were added to the 2 ml sample, followed by the addition of concentrated sulfuric acid drop by drop. The addition caused a change in color to either pink scarlet or crimson red.

Zinc chloride reduction assay: An admixture of zinc dust with strong hydrochloric acid was stirred into the 2 ml sample, and over time, a red color appeared.

Alkaline indicator test: Adding sodium hydroxide solution to the 2 ml sample led to a color change, turning yellow or red.

2.5.4. Tanninidentification:

Gelatinprocedure:A 0.01% gelatin solution mixed with 10% sodium chloride was added to the 2 ml sample, resulting in a white solid settling ppt.

Ferricchloridereaction:When ferric chloride was introduced to the 2ml sample, it resulted in a blue-green color.



Figure 5: Tannin Identification

2.5.5. Proteins and amino acid analysis:

Millon's indicator: The addition of Millon's reagent to the 2ml sample initially produced a white precipitate, which turned red when heated.

Ninhydrin assay: Combining the 2 ml sample with a 0.2% ninhydrin solution and heating produced plum violet color, indicating the existence of proteins as well as amino acids.

2.5.6. Fattests:

Stain test: A little portion of sample being placed intermediary in filter paper sheets, and any oil stain obtained on filter sheet exclaimed the existence of non-volatile oils.

Saponification analysis: Adding small quantity of 0.5 Normal potassium hydroxide (alcoholic) to the sample, phenolphthalein a small drop, and heating it with water bath for almost 2 hours resulted in the development of soap or the near complete annulment of the base, indicating the existence of fats and non-volatile oils.

2.5.7. Sterol screening:

Liebermann-Burchard method: By adding 3-4 drops of acetic anhydride, boiling the mixture, cooling, and then adding 3 drops of conc. sulphuric acid with the help of a dropper, the brown ring formed at the confluence, indicating the existence of steroids.

2.5.8. Triterpenoid detection:

Salkowski assay: When a 2 ml portion of chloroform was mixed with some drops of concentrated H₂SO₄ and thoroughly shaken, reddish-brown color in the underneath bands suggested the existence of steroids, while yellow color in the upper band affirmed the existence of triterpenoids.

2.5.9. Foaming Index Visualization:

The Visualization of Foaming Index assesses the presence of saponins if any, which are high molecular weight phytoconstituents known for their detergent properties. Some medicinal plants are having saponins that can produce stable foam, when their hydrous decoction is agitated. The lathering capacity of the plant material and its extracts is quantified using a foaming index. A precise weight of 1 gram of coarse powder of *Rhaphidophora aurea* is used for this measurement.



Figure 5: Foaming Index

2.5.10. Swelling Index:

Some herbal drugs possess therapeutic or pharmaceutical value due to their swelling properties, particularly gums and those with mucilage, pectin, or hemicelluloses. The swelling index was measured in ml. Accurately weighted 1g of *Rhaphidophora aurea* powder was transferred to a 50ml measuring cylinder. 25 ml of water was added successively, and mixture was shaken vigorously at every 10 minutes for 1 hour. Allow the mixture to stand for 4 hours at room temperature. volume in ml was measured that occupied by the plant material, including any sticky mucilage.



Figure 6: Swelling Index

2.6. Sterility test of the plant extracts:

The plant's methanolic extract underwent testing for microbial growth. This involved inoculating 0.5ml of each sample on sterile Mueller Hinton Agar and then incubating them over 37°C for almost 24 hours. Fungal growth was assessed on Sabouraud Dextrose Agar over a 4-day period at 25°C. The plates were monitored for any type of bacterial or fungal growth. If no sign of growth was observed in the extracts after incubation, it suggests sterility and was further assessed for antimicrobial activity following CLSI guidelines.



Figure 7: Sterility Test of Plant Extracts

3. Results:

3.1. Macroscopic and Microscopic Evaluation

3.1.1 Macroscopic Analysis:

The macroscopic evaluation of *Rhaphidophora aurea* leaves provided distinct morphological characteristics, aiding in the authentication and identification of the plant material. The observed characteristics are summarized as follows:

- **Leaf Shape:** Broadly ovate to heart-shaped with an acuminate apex and entire margins.
- **Size:** Average leaf size measured 10-15 cm in length and 6-10 cm in width.
- **Color:** Fresh leaves exhibited a vibrant green color with yellow variegation, while dried leaves appeared pale green to brownish-green.
- **Texture:** The leaves were smooth, glossy, and leathery with a slightly waxy cuticle.
- **Venation Pattern:** Pinnate venation with a prominent midrib and secondary veins running parallel to the leaf margin.
- **Surface Features:**
 - **Adaxial (Upper) Surface:** Shiny, with a slightly thickened cuticle.
 - **Abaxial (Lower) Surface:** Slightly rough with visible stomatal openings and scattered trichomes.

These macroscopic features correlate with the general characteristics of the Araceae family, facilitating plant identification (Kokate et al., 2019).



Figure 8: Macroscopic Features

3.1.2 Microscopic Analysis:

The transverse section of *Rhaphidophora aurea* leaf revealed the following microscopic structures:

Epidermis: The leaf exhibited a single-layered epidermis covered with a thick cuticle. The adaxial (upper) epidermis contained compactly arranged cells with less stomata. The abaxial (lower) epidermis had a higher density of paracytic stomata, characteristic of the Araceae family.




Figure 9: Microscopic Image of epidermis

Trichomes:Non-glandular unicellular trichomes were present on the lower epidermis.The trichomes were thin-walled and elongated, aiding in protection and reducing water loss.



Figure 10: Microscopic Image of Trichome

Mesophyll:The mesophyll was differentiated into palisade and spongy parenchyma.The palisade parenchyma was columnar, tightly packed, and contained abundant chloroplasts for photosynthesis.The spongy parenchyma consisted of loosely arranged cells with prominent intercellular spaces, assisting in gaseous exchange.




Figure 11: Microscopic Image of Mesophyll

Vascular Bundles:The vascular system was collateral, with xylem facing the adaxial surface and phloem toward the abaxial side.Xylem vessels were lignified, providing mechanical support.Phloem was composed of sieve elements and companion cells, facilitating nutrient transport.The vascular bundles were surrounded by a bundle sheath of parenchymatous cells.




Figure 12: Microscopic Image of Vascular Bundles

B. Powder Microscopy

The powdered leaf sample exhibited various diagnostic characters under microscopic examination:

- **Epidermal Cells:** Polygonal, with thickened walls and a cuticular layer.
- **Stomata:** Predominantly paracytic stomata, commonly found in monocot plants of the Araceae family.
- **Trichomes:** Few scattered unicellular trichomes, observed on the abaxial epidermis.
- **Lignified Xylem Vessels:** Spiral and annular thickening patterns were visible with phloroglucinol-HCl staining, indicating lignification.
- **Calcium Oxalate Crystals:** Prismatic and rosette-shaped calcium oxalate crystals were identified, confirming the characteristic feature of Araceae family plants.
- **Starch Grains:** Oval to polygonal starch grains, confirmed by iodine-potassium iodide (I₂KI) staining, indicating carbohydrate storage.




Figure 13: Powder Microscopic images

3.2. SEM Analysis:

The SEM analysis of *Rhaphidophora aurea* leaf powder revealed key micromorphological features critical for plant identification and quality control. The presence of paracytic stomata, unicellular trichomes, vascular elements with spiral xylem thickening, and calcium oxalate crystals strongly supports the plant's pharmacognostical identity. The observations align with previous reports on the Araceae family, confirming the diagnostic characteristics essential for standardization and authentication of *Rhaphidophora aurea* in herbal formulations.




Figure 14: SEM images *Rhaphidophora aurea* leaf

3.3. Physicochemical Evaluation:

The physicochemical evaluation of *Rhaphidophora aurea* leaf powder was conducted to establish **standardization parameters** and ensure **quality control** of the plant material. The results obtained from various physicochemical tests, including **moisture content, ash values, and extractive values**, are summarized below in Table 1.

Table 01: Ash value of *Rhaphidophora aurea* by soxhlet extraction method

Sr.no.	Physical contents	Value(%w/w)
1	Moisture Content	5.42 ± 0.13
2	Total Ash value	9.68 ± 0.21
3	Acid insoluble ash	1.74 ± 0.09
4	Water soluble ash	3.87 ± 0.15

The moisture content of *Rhaphidophora aurea* was determined using the Loss on Drying (LOD) method at 105°C for one hour. The low moisture content suggests that the plant material has a longer shelf life with minimal microbial contamination risk. The total ash content, which represents the total amount of inorganic matter present in the sample after complete incineration. A higher total ash value may indicate the presence of minerals, but excessive ash content could suggest contamination with extraneous matter such as soil, sand, or inorganic adulterants. The water-soluble ash content, representing the portion of the total ash that is soluble in water. Water-soluble ash indicates the amount of bioavailable inorganic components, such as potassium, calcium, and sodium salts, which may contribute to the plant's medicinal properties. A low water-soluble ash value suggests limited mineral content in the plant, confirming minimal contamination by water-soluble extraneous matter. The acid-insoluble ash value, which indicates the amount of siliceous matter. A low acid-insoluble ash value suggests minimal contamination with siliceous matter.

3.4. Extractive value:

The extractive values of *Rhaphidophora aurea* leaf powder were determined using alcohol and water as solvents to assess the presence of soluble bioactive compounds. These values provide insights into the quantity of phytochemicals extracted from the plant material and help in standardization and quality control as shown in table 2.

Table 02: The Extractive values obtained from the leaf powder of *Rhaphidophora aurea* by soxhlet extraction method:

Sr.no.	Extract	Values(%w/w) by soxhlet Extraction
1	Alcohol-Soluble Extract	11.32 ± 0.18
2	Water-Soluble Extract	9.54 ± 0.20




Figure 15: Physicochemical Parameters Analysis

3.5. Phytochemical Analysis:

Chemical Test	Inference	Methanol (80%)	Chloroform	Petroleum Ether
Mayer's test	Cream coloured	+	+	+
Dragendroff's test	Reddish brown precipitate	+	+	+
Wagner's test	Reddish-brown	+	+	+
Hager's test	Yellow coloured precipitate	+	+	+
Fehling's test	Brick red precipitate	-	-	-
Legal's test	Blood red colour	-	-	-
Shinoda test	Pink scarlet	+	+	+
Zinc Chloride reduction test	Red colour	+	+	+
Alkaline reagent test	Yellow colour	+	+	+
Gelatin test	White precipitate	+	+	+
Ferric Chloride test	Blue green colour	+	+	+
Millon's test	No White sediment	-	-	-
Ninhydrin test	Violet Colour	-	-	-
Liebermann-Burchard test	Brown Ring Appears	+	+	+
Salkowski test	No colour	-	-	-

The presence of alkaloids was confirmed in all three solvent extracts (Methanol, Chloroform, and Petroleum

Ether) through Mayer's, Dragendroff's, Wagner's, and Hager's tests. Flavonoids were detected in all extracts using Shinoda, Zinc Chloride Reduction, and Alkaline reagent tests. Tannins were confirmed through Gelatin and Ferric Chloride tests. Sterols were present in all extracts as indicated by the Liebermann-Burchard test. Glycosides were absent in all extracts. Terpenoids were not detected in any of the extracts. Proteins and amino acids (Millon's and Ninhydrin tests) were absent in all extracts.

4. Discussion:

The present study extensively evaluates the pharmacognostical, physicochemical, and phytochemical attributes of *Rhaphidophora aurea* (Money Plant), an ornamental plant that has demonstrated potential medicinal properties. The results contribute to scientific standardization, authentication, and possible therapeutic applications. Each analytical aspect of the study was designed to ensure the quality, purity, and botanical reliability of the plant material, facilitating its future inclusion in herbal pharmacopoeias.

The macroscopic examination of *Rhaphidophora aurea* confirmed its characteristic ovate to heart-shaped leaves, glossy surface, prominent pinnate venation, and thick leathery cuticle. These features align with those observed in members of the Araceae family, reinforcing its botanical classification. The leaf dimensions, surface texture, and color variations provide vital identification markers for plant authentication and differentiation from potential adulterants.

The microscopic analysis, including transverse sections of the leaf and powder microscopy, provided insights into the plant's anatomical structure. The epidermis displayed a single-layered arrangement with a thick cuticle, which is known to aid in water retention and protection against environmental stress. The presence of paracytic stomata, a distinguishing feature of the Araceae family, was confirmed primarily on the abaxial surface, supporting gas exchange and transpiration processes.

Furthermore, the vascular bundles were found to be collateral, with xylem vessels exhibiting lignified spiral thickening. This characteristic ensures the plant's structural integrity and efficient transport of nutrients and water. The presence of unicellular non-glandular trichomes, particularly on the lower epidermis, suggests that the plant possesses natural defense mechanisms against herbivory and environmental stress. Additionally, the identification of calcium oxalate crystals, starch grains, and mucilage cells further strengthens its botanical characterization. These findings establish the plant's micromorphological uniqueness and serve as diagnostic markers for its pharmacognostical standardization.

The physicochemical evaluation of *Rhaphidophora aurea* was performed to establish quality control parameters, ensuring the plant's purity and safety for potential medicinal applications. The moisture content was determined to be $5.42 \pm 0.13\%$, which falls within acceptable limits as per World Health Organization (WHO) standards for herbal drugs. A moisture content below 10% is considered suitable for long-term storage since it minimizes the risk of fungal and microbial contamination. This result confirms that *Rhaphidophora aurea* is relatively stable and can be preserved without significant degradation over time. The total ash value was found to be $9.68 \pm 0.21\%$, indicating a moderate presence of inorganic residues. The water-soluble ash value ($3.87 \pm 0.15\%$) signifies the proportion of bioavailable minerals such as potassium, sodium, and calcium, which may contribute to the plant's nutritional and therapeutic properties. Additionally, the acid-insoluble ash value ($1.74 \pm 0.09\%$) suggests minimal contamination from siliceous matter such as soil, sand, or dust. These values confirm the plant material's purity and suitability for medicinal use, ensuring compliance with pharmacopoeial quality standards. The extractive values help determine the amount of bioactive compounds present in the plant material. The alcohol-soluble extractive value was found to be $11.32 \pm 0.18\%$, indicating a high presence of ethanol-soluble phytoconstituents such as flavonoids, alkaloids, and steroids. These compounds are known for their antioxidant, anti-inflammatory, and antimicrobial properties, making alcohol-based extracts of *Rhaphidophora aurea* valuable in herbal medicine. Similarly, the water-soluble extractive value ($9.54 \pm 0.20\%$) indicates the presence of hydrophilic compounds such as glycosides, tannins, and carbohydrates, which are often responsible for the astringent, antimicrobial, and immunomodulatory effects of medicinal plants. The relatively high extractive values suggest that *Rhaphidophora aurea* contains a significant proportion of bioactive compounds, making it suitable for both aqueous and ethanol-based herbal preparations.

The phytochemical analysis of *Rhaphidophora aurea* confirmed the presence of various secondary metabolites, which contribute to its medicinal potential. All four alkaloid detection assays (Mayer's, Wagner's, Dragendroff's, and Hager's tests) confirmed the presence of alkaloids in methanol, chloroform, and petroleum ether extracts. Alkaloids are well-known for their analgesic, neuroprotective, and antimicrobial properties, indicating that *Rhaphidophora aurea* may have pharmacological relevance in these domains. The Shinoda test, Zinc chloride reduction test, and Alkaline reagent test confirmed the presence of flavonoids in all extracts. Flavonoids are potent antioxidants known for their anti-inflammatory, anti-cancer, and cardiovascular benefits. This result strengthens the argument for *Rhaphidophora aurea* as a potential source of natural antioxidant

compounds. Tannins were detected in all extracts through Gelatin and Ferric chloride tests, suggesting astringent and antimicrobial activity. Tannins are commonly used for wound healing, gastrointestinal protection, and antimicrobial treatments. Sterols were confirmed in all extracts via the Liebermann-Burchard test, indicating their membrane-stabilizing and anti-inflammatory properties. However, terpenoids were not detected, suggesting a lesser role in the plant's medicinal profile. Glycosides and proteins were not detected in any of the extracts, indicating that these compounds do not contribute significantly to the plant's therapeutic properties.

The **SEM analysis** provided insights into the **micromorphological features** of *Rhaphidophora aurea*, confirming:

- **Paracytic stomata with distinct guard cells**, aiding in respiration.
- **Unicellular trichomes**, which protect against external stressors.
- **Well-developed vascular bundles**, with **spiral xylem thickening**, supporting efficient nutrient transport.
- **Calcium oxalate crystals**, which serve as diagnostic markers for the **Araceae family**.

These observations validate the **anatomical identity of *Rhaphidophora aurea***, strengthening its pharmacognostical standardization.

The results of this study establish pharmacognostical markers that can be used to authenticate and standardize *Rhaphidophora aurea* for medicinal use. The presence of flavonoids, tannins, sterols, and alkaloids indicates its therapeutic potential in herbal formulations. Furthermore, the low moisture content and optimal ash values ensure its suitability for long-term storage and formulation stability.

While *Rhaphidophora aurea* has been widely recognized for its air-purifying benefits, this study highlights its medicinal attributes, paving the way for its inclusion in traditional and modern herbal pharmacopoeias.

5. Conclusion:

The comprehensive evaluation of *Rhaphidophora aurea* confirms its botanical identity, purity, and bioactive potential. The study establishes quality control parameters, supporting its safe and effective use in herbal medicine. Further studies should focus on quantifying specific bioactive compounds and evaluating pharmacological activities to unlock the full medicinal potential of this plant.

Reference:

1. Kriti Saxena, Rajat Yadav and Dharmendra Solanki, *Rhaphidophora aurea*: A Review on Phytotherapeutic and Ethnopharmacological Attributes. International Journal of Pharmaceutical Science Review and Research, 69(1), 2021, Pages: 236-247.
2. Poole RT, Chase AR, Osborne LS, CFREC-A Foliage Plant Research Note RH- 1991-29, journal of Institute of Food and Agricultural Sciences, Central Florida Research and Education Center, Apopka; 2018; 5:44-49.
3. Liliwirianis N, Wan Z, Wan MZ, Kamaruzaman J, Screening of *Epipremnum* Sp., *Syngramma alysmifolia*, *Thotea* Sp. and *Smilax* Sp. For Antimicrobial Activity; journal of World Applied Sciences, 2010; 8 (7): 889-891.
4. Meshram A, Srivastava N. Molecular and physiological role of *Epipremnum aureum*. International Journal of Green Pharmacy 2014; 8: 73-76.
5. Bakhuizen VD, Brink RC, Are *Epipremnum* Schott, *Rhaphidophora* Hassk., and *Monstera Adanson* congeneric.
6. Blume A Supplement, International Journal of Green Pharmacy, 1958; 4: 91-92.
7. Jed W F, Moringa SCD, A Review of the medical evidence for its nutritional, therapeutic, and prophylactic properties, Trees for life Journal, Part 1: 42-54
8. Ahmad I, Beg AZ, Antimicrobial and phytochemical studies on 45 Indian medicinal plants against multi-drug resistant human pathogens, Journal of Ethnopharmacology, 2001; 74: 113-123.
9. Ikram M, Inamul H, Screening of medicinal plants for antimicrobial activities, Journal of Pharmaceutical Research & Clinical Practice, 1984; 10: 62-64.
10. Cowan MM, Plant products as antimicrobial agents, journal of Clinical Microbiological Review, 1999; 12: 564-582.
11. Parulpriya P, Lalitha P, Phytochemical constituents and antimicrobial activity of the aerial roots of *Rhaphidophora aurea* at wineedover *Areca catechu*, National conference on natural product research: for health and biopharma industry, 2011; 12: 24
12. Arathi KA, Lalitha P, Jayanthi P, Shubashini K S, Phytochemical investigation of the Aerial roots of *Pothos aurea* (Linden ex Andre), report of 3rd Indo- Korean joint seminar on medicinal plant research,

- 2010: 53.
13. Lalitha P, Arathi KA, Shubashini K S, Hemalatha S, Jayanthi P, Antimicrobial activity and Phytochemical screening on an ornamental foliage plant *Pothos aurea*, Asian journal of Pharmaceutical and clinical research, 2010; 1(2):1-07.
 14. Tani A, Nicholas H C, Uptake of aldehydes and ketones at typical indoor concentrations by houseplants, journal of Environmental Science and Technology, 2009; 43 (21):8338–8343.
 15. Parulpriya P, Lalitha P, Efficiency of different solvent extraction methods in the isolation of bioactive principles from the ethyl acetate extract of *Rhaphidophora aurea*, International conference in “Global competitiveness in biology, chemistry and environment for a better tomorrow, 2011; 10: 48
 16. Meshram A, Srivastava N. ”epipremnum aureum (jade pothos): a multipurpose plant with its medicinal and pharmacological properties” Journal of Critical Review; vol 2 (issue 2); 2015: 45-46.
 17. Abhigyan G, Naga K R, Fathima B, Agarwal P, Anti diabetic activity of *Epipremnum aureum* in normal and alloxan induced diabetic Rats; Asian journal of Pharmaceutical and clinical research ; Vol.9 ; issue 4; 2016: 54-57.
 18. Srivastava N, Shwarupa S, Bhagyawant SS, Comparative study on the anti-termite, antimicrobial and antioxidant activity of leaf and root extracts of *Pothos aurea* (*Epipremnum aureum* L.), Journal of Pharmaceutical Research and Clinical Practice 2011; 1:111.
 19. Yang H, Liu Y. Phytoremediation on air pollution In: The impact of air pollution on health, economy, environment and agricultural, international journal of environmental technology; 2011; Vol 1: 281-94.
 20. Douglass F. Indoor air quality-planting healthier indoor air, News forum Environmental Health Perspectives, 2011; 119: A426-427.
 21. Lan TM, Sulaiman SF, Najimudin N, Muhammad T ST, Anticancer medicinal plant *Epipremnum pinnatum* chloroform extracts elicited both apoptotic and non-apoptotic cell deaths in T-47D mammary carcinoma cells, KMITL Science and Technology Journal 2007; 7: 24-43.
 22. Palmer M, Betz JM. Plants. In: Goldfrank LR, Flomenbaum NE, Hoffman RS, Howland MA, Lewin NA, Nelson L, editors. Goldfrank Toxicologic Emergencies. 8; 2006: 1577-602.
 23. Meshram A, Srivastava N. *In vitro* antitermite activity of alkaloids from *Epipremnum aureum* (Linden and Andre) Bunting (Araceae) against Indian white termite *Odontotermes obesus*. Asian Journal of Pharmaceutical Technology and Innovation, 2015; 3(10):27-31.
 24. Weidner M, Silva JAT, Potential and limitations of ornamental plants for indoor air purification, Journal of Floriculture, ornamental and plant biotechnology, Advances and Topical Issues, 2006; 4: 54-63.
 25. Liu ZC, Lu SM, Liu CB, Zhao YF, Wang HL, Li JM, *et al.* Simultaneous determination of nicotine and cotinine in leaves of plants in ETS by UPLC, the journal of Chemical Engineering 2008; 12:43-45.
 26. Weidner M, Martins R, Muller A, Simon J, Schmitz H. Uptake, transport and accumulation of nicotine by the Golden pothos (*Epipremnum aureum*): the central role of root pressure. J Plant Physiology 2005; 162: 139-50.
 27. Zheng-cong IU, She-ming LU, Chun-bo LIU, Yun-fang ZHAO, Hai-li WANG, Jinming LI, Ming-ming MIAO, Simultaneous determination of nicotine and cotinine in leaves of plants in ETS by UPLC, Chemical engineer, 2008; 12: 33.
 28. Manfred Weidner, Ralf Martins, Andrea Muller, Judith Simon, Heribert Schmitz Uptake, transport and accumulation of nicotine by the *Golden potho* (*Epipremnum aureum*): the central role of root pressure, Journal of plant physiology, 2019; 162: 139-150.
 29. Normon DJ, Yuen JMF, A distinct pathotype of *Ralstonia* (*Pseudomonas*) *solanacearum* race 1, biovar 1 entering Florida in *Pothos* (*Epipremnum aureum*) cuttings, Canadian journal of plant pathology, 2015; 20: 171-175.
 30. Heather L Papinchak, Jay Holcombe, Teodora Orendovici Best, Dennis R Decoteau, Effectiveness of houseplants in reducing the indoor air pollutant ozone, journal of Horticulture Technology, 2009; 19(2): 286-290.
 31. Yuichi T, Matsuzaki T, Yuya T, Isolation and characterization of formaldehyde-responsive genes from Golden pothos (*Epipremnum aureum*), journal of Plant biotechnology, 2010; 27:325–331.
 32. Kil MJ, Kim KJ, Kim HH, Lim YW, Effects of growing media and exposure frequency on the volatile formaldehyde removal in potted *Epipremnum aureum*, Korean journal of Horticultural Science and Technology, 2008; 26(3): 325-330.
 33. Jeong SJ, Kim WS, Lee DW, Kim HD, Kim KJ, Yoo EH, Cho JG, Evaluation of selected foliage plants for improvement of indoor humidity, horticulture, environment, and biotechnology, Korean journal of Horticultural Science and Technology, 2008; 49(6):439-446.

34. Park J.S, Masahisa Ishii, Limi O, Sadanori S, Kazuhiro Fujiwara, Effects of ozonated water-fog cooling on ozone gas concentrations and population densities of airborne bacteria and fungus on plant leaves in a naturally ventilated greenhouse, *The Japan Agricultural Research Quarterly*, 2008; 42(4): 291–297.
35. Tarran J, Torpy F, Burchett M, Use of living pot-plants to clean indoor air – research review, *Proceedings of sixth international conference on indoor air quality, Ventilation & energy conservation in buildings – Sustainable built environment*, 2007; Volume III: 249-256.
36. Orlikowski LB, Trzewik A, Wiejacha K, Phytophthor tropical ison *Hederahelix* and *Epipremnum aureum* in Polish greenhouses, *Journal of Communicative Agriculture and Applied Biology*, 2019; 5: 56-67.
37. Lalitha P, Arathi KA, Sripathi SK, Hemalatha S, Jayanthi P. Antimicrobial activity and phytochemical screening of an ornamental foliage plant, *Pothos aurea* (Lindenex Andre). *Alfa Universal: An International Journal of Chemistry*, 2010; 1(2): 63-71.
38. Sawada Ayako, Kuroda Hiroyuki, Yoshida Taketoshi, Oyabu Takashi, Takenaka Kozaburo, Purification effects of Golden pothos and peace lily for indoor air- pollutants and its application to a real environment, *Papers of technical meeting on chemical Sensor, journal of Institute of electrical engineers, Japan*, 2004; 04(1)-13: 57-62.
39. Sriussadaporn C, Yamamoto K, Fukushi K, Simazaki D, Comparison of DNA damage detected by plant comet assay in roadside and non-roadside environments, *journal of Mutational Research*, 2003; 541(1-2): 31-44.
40. Heribert S, Hilgers U, Weidner M, Assimilation and metabolism of formaldehyde by leaves appear unlikely to be of value for indoor air purification, *journal of Research of new phytologists*, 2000; 147(2): 307–315.
41. Paulsen E, Stahl S P, Andersen KE, Immediate skin and mucosal symptoms from pot plants and vegetables in gardeners and greenhouse workers, *Contact dermatitis, Journal of Health sciences and plants*. 1998; 39(4): 166– 170.
42. Nihal C R, John W, David W M R, Use of antitranspirants under low-light environments to control transpiration of *Epipremnum aureum* leaves, *Scientia Horticulturae*, 43; (3-4): 307-312.
43. Seong H P, Randomized clinical trials evaluating therapeutic influences of ornamental indoor plants in hospital rooms on health outcomes of patients recovering from surgery, a dissertation submitted in partial fulfillment of the requirements for the degree. Doctor of philosophy, Department of Horticulture, Forestry and Recreation Resources, College of Agriculture, *journal of Green Pharmacy*, 2006; 4: 21-32.
44. Khayyat M, Nazari F, Salehi H, Effects of Different Pot Mixtures on *Pothos Epipremnum aureum* Linden. And Andre ‘Golden pothos ’ Growth and Development, *American-Eurasian Journal of Agricultural & Environmental Science*, 2007; 2 (4): 341-348.
45. Luping QU, Jianjun C, Richard J, Henny, Yingfeng H, Russell D, Caldwell C A, Robinson, Thidiazuron promotes adventitious shoot regeneration from *Pothos* (*Epipremnum aureum*) leaf and petiole explants, *international Journal of In Vitro Cellular Development and Biology of Plant*, 2001; 38: 268–271.
46. Huxley A, Griffiths M, Levy M, Griffiths M. *Index of garden plants, The New Horticulture Society Dictionary of Gardening*, 4th ed. Vol. 1. London: Macmillan; 1994: 3353.
47. Ogundipe O, Akinbiyi O, Moody JO. Antimicrobial activity of selected ornamental plants. *Niger Journal of National Products of Medicines*, 1998; 2: 46-8.
48. Yang H, Liu Y, Phytoremediation on air pollution, the impact of air pollution on health, economy, environment and agricultural sources. *International journal of technology*; 2011: 281-94.
49. Douglass F, Indoor air quality planting healthier indoor air, *News forum Environmental and Health Perspectives* 2011; 119: A426-A427.
50. Xueng Z, Wang L, Hou H. Formaldehyde removal by potted plant-soil systems. *Journal of Hazardous Material*, 2011; 192: 314–318.
51. Wolverton BC, Johnson A, Bounds K. Interior landscape plants for indoor air pollution abatement, *Final Report September NASA Stennis Space Center, MS*; 1989: A45-A 46
52. Davanzo F, Miaglia S, Perego S, Assisi F, Bissoli M, Borghini R, *et al.* Plant poisoning: Increasing relevance, a problem of public health and education. North-Western Italy, *Journal of Pharmacy Scientific Research*, 2011; 3(7): 1338-43.
53. Palmer M, Betz JM, Goldfrank LR, Flomenbaum NE, Hoffman RS, Howland MA, Lewin NA, Nelson L. editors. *Goldfrank’s journal of Toxicological Emergencies*. 8th ed.; 2006: 1577-602.
54. Hung CY, Sun YH, Chen J, Darlington DE, Williams A, Burkey KO, *et al.* Identification of a Mg protoporphyrin IX monomethyl ester cyclase homologue, EaZIP, differentially expressed in variegated *Epipremnum aureum* ‘Golden Pothos’ is achieved through a unique method of comparative

- study using tissue regenerated plants, *Journal of Experimental Botany*, 2010; 61:1483-93.
56. Desai TS. Studies on thermoluminescence, delayed light emission and oxygen evolution from photosynthetic materials: UV effects, *Journal of Photosynthetic Research*, 1990; 25:17-24.
 57. Arathi KA, Lalitha P, Jayanthi P, Shubashini K Sripathi, Phytochemical investigation of the Aerial roots of *Pothosaurea* (Linden ex Andre), journal 3rd Indo Korean joint seminar on medicinal plant research, 2010; 5: 53-54
 58. Patel, Roshan, Diuretic activity of leaves of *Plectranthus ambionicum* spring in male albinorats, *the journal of Pharmacognostical research*, 2010; 2.2: 86.
 59. Dev, Protopaditya, Neuropharmacological properties of *Milkiania scandens* (linn) Willd (astraceae), *Journal of advanced Pharmaceutical technology and research*, 2011; 214: 255
 60. Gahlot K, Abid M, Sharma A, Pharmacological evaluation of *Gelenium sempervirens* roots of CNS depressant activity, *International Journal of Pharmaceutival research*, 2011; 3.2: 693-697
 61. Koga K, Iwasaki Y. Psychological and physiological effect in humans of touching plant foliage-using the semantic differential method and cerebral activity as Indicators, *Journals of Physiology and Anthropology*, 2013; 32: 7.
 62. Wan Z, Wan MZ, Liliwirianis N, Kamaruzaman J, Screening of *Epipremnum* Sp., *Syngramma alysmifolia*, *Thote a* Sp. and *Smilax* Sp. For Antimicrobial Activity, *journal of World Applied Sciences*, 2010; 8 (7): 889-891.
 63. Sriwasatava N, Shwarupa S, Bhagyawant S.S., Comparative Study on the Anti-Termite, Antimicrobial and Antioxidant Activity of Leaf and Root Extracts of *Pothosaurea* (*Epipremnum aureum* L.), *Journal of Pharmaceutival Research & Clinical Practice*, Apr-June 2011; 1(2): 1-11.
 64. The Removal of Fluoride from the Water Using *Epipremnum aureum* (Money Plant) - Scientific Figure on ResearchGate. Available from: https://www.researchgate.net/figure/SEM-image-of-the-surface-of-the-money-plant-dry-leaves-powder-MLDP_fig1_373714251 [accessed 22 Jul 2024].